# Comparison of Decolorization of Methylene Blue Aqueous Solutions Irradiated with Atmospheric-Pressure Plasma Jet in Air and in Argon Gas

by

Hiroshi KUWAHATA<sup>\*1</sup>, Masato UCHIDA<sup>\*2</sup> and Yoshiki ODA<sup>\*3</sup>

(Received on Mar. 30, 2018 and accepted on May 10, 2018)

#### Abstract

The decolorization of a methylene blue aqueous solution irradiated with an atmospheric-pressure argon (Ar) plasma jet in air was compared with that in Ar gas. The time required for decolorization of the methylene blue aqueous solution irradiated with a plasma jet in Ar gas was 10 min, which was half that required for decolorization in air. We suggest that this is caused by a high current of 60.8 mA flowing during plasma jet irradiation in Ar gas, which is ~1.6-fold higher than that in air. In addition, in the case of the plasma jet irradiation in Ar gas, neither nitrate ions (NO<sub>3</sub><sup>-</sup>) nor nitrite ions (NO<sub>2</sub><sup>-</sup>) were generated, but only hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) was generated, unlike the case of plasma jet irradiation in air. The results obtained in this study show that harmful organic compounds in wastewater can be decomposed quickly without generating nitrogen oxide ions (NO<sub>x</sub><sup>-</sup>) by irradiating the solution with an atmospheric-pressure Ar plasma jet in Ar gas, compared with the case of irradiation in air.

Keywords: Atmospheric-pressure plasma, Methylene blue aqueous solution, Decolorization

### 1. Introduction

Research on wastewater treatment using discharge and plasma has been attracting attention: for example, wastewater treatment using surface pulse discharge <sup>1-5)</sup>, surface creeping discharge <sup>6)</sup>, surface atmospheric-pressure plasma <sup>7-10)</sup>, underwater pulse discharge <sup>11-13)</sup>, and underwater discharge inside bubbles <sup>14-17)</sup> have been reported thus far. In these reports, methylene blue ( $C_{16}H_{18}CIN_3S$ ), a blue organic dye, has been frequently used as an indicator of decomposition <sup>4, 5, 7, 9, 15-17)</sup>. Hydroxyl (OH) radicals generated by dissociation of water (H<sub>2</sub>O) molecules as a result of collisions of high-energy electrons in the discharge and plasma with H<sub>2</sub>O molecules are associated with the decomposition of organic compounds. OH radicals are reactive oxygen species (ROS), have a high oxidation potential (2.80 V) <sup>18-20)</sup>, and oxidatively decompose organic compounds.

We previously reported that a methylene blue aqueous solution was decolorized when it was irradiated with an atmospheric-pressure argon (Ar) plasma jet in air for 20 min <sup>21)</sup>. We also found, on the basis of the results of nuclear magnetic resonance (NMR) and electrospray ionization mass spectroscopy (ESI-MS), that the decolorization is associated with the decomposition of the phenothiazine ( $C_{13}H_9NS$ ) structure accompanied by the dissociation of a methyl (CH<sub>3</sub>) group from a side chain of the methylene blue molecule <sup>22)</sup>.

In the previous articles, we reported that when distilled water is irradiated with an Ar plasma jet, nitrate ions  $(NO_3^-)$  are generated in distilled water and the absorption spectrum has a peak at a wavelength of ~200 nm <sup>23)</sup>. This  $NO_3^-$  is considered to be generated as follows: high-energy electrons in plasma collide with nitrogen  $(N_2)$  and oxygen  $(O_2)$  molecules in air, inducing dissociation to form nitrogen (N) and oxygen (O) atoms. Then, nitrogen oxide ion  $(NO_x^-)$  species are generated by the bonding of N and O atoms. When  $NO_x^-$  enters distilled water, nitric acid  $(HNO_3)$  is generated. Finally,  $HNO_3$  is ionized to generate  $NO_3^-$ . We also found that the peak at a wavelength of ~200 nm

<sup>\*1</sup> Associate Professor, Department of Electrical and Electronic Engineering

<sup>\*2</sup> Graduate Student, Course of Electrical and Electronic System

<sup>\*3</sup> Technician, Technology Joint Management Office

corresponds to the presence of hydrogen peroxide  $(H_2O_2)$  and a small amount of nitrite ions  $(NO_2^-)$ , from the results of high-performance liquid chromatography (HPLC)<sup>24)</sup>. From these results, we considered that  $NO_3^-$ ,  $NO_2^-$ , and  $H_2O_2$  are generated in a methylene blue aqueous solution irradiated with an Ar plasma jet; consequently, a peak at a wavelength of ~200 nm should exist in the absorption spectra.

In this study, a methylene blue aqueous solution was irradiated with an Ar plasma jet and absorption spectroscopy was carried out. If there is a peak at a wavelength of  $\sim 200$ nm in absorption spectra, then  $NO_3^-$ ,  $NO_2^-$ , and  $H_2O_2$  are present in the methylene blue aqueous solution. HPLC was carried out to confirm the generation of NO<sub>3</sub>, NO<sub>2</sub>, and  $H_2O_2$ . If  $NO_3^-$  and  $NO_2^-$  are generated as a result of collisions between high-energy electrons in plasma with N2 and  $O_2$  molecules in air,  $NO_3^-$  and  $NO_2^-$  should not be generated in the absence of the contact between plasma and air during irradiation. On the basis of this idea, a methylene blue aqueous solution was irradiated with a plasma jet in Ar gas, instead of air, and absorption spectroscopy and HPLC were carried out on the resultant solutions. Emission spectroscopy of the Ar plasma jet was also carried out to confirm the absence of N2 molecules during plasma irradiation in Ar gas. The times required for decolorization of the solution irradiated with a plasma jet in Ar gas and air were different, and the currents flowing during plasma jet irradiation in Ar gas and in air were compared.

# 2. Experimental Methods

Figure 1 shows a schematic of the experimental setup used in this study. In the unit used to generate the plasma jet, a copper tube (inner diameter, 4 mm; outer diameter, 6 mm), as the discharge electrode, was inserted into a dielectric quartz tube (length, 50 mm;inner diameter, 6 mm; outer diameter, 8 mm) around which copper foil (thickness, 0.05 mm; width, 10 mm) was wrapped as the ground electrode <sup>25)</sup>. A high AC voltage was applied to induce dielectric barrier discharge inside the quartz tube between the electrodes and to convert the Ar gas to plasma that was then ejected into the air in the form of a jet. A high-voltage power supply (LHV-10AC, Logy Electric Co., Ltd.) was used to generate the Ar plasma jet (frequency, 10 kHz; applied voltage, 10 kV; Ar gas flow rate, 10 L/min). A methylene blue aqueous solution [10 mL of a 10 mg/L (10 ppm) solution] in a glass Petri dish (diameter, 60 mm) was irradiated with the Ar plasma jet in air for 1-20min. The distance from the end of the quartz tube to the solution (irradiation distance) was 15 mm. Plasma jet irradiation in Ar gas was carried out after an acrylic resin container  $(80 \times 80 \times 30 \text{ mm}^3)$  was filled with



Fig. 1 Schematic of the experimental setup.



Fig. 2 Circuit used to measure current.

Ar gas. The voltage and current were measured using a high-voltage probe (P6015A, Tektronix, Inc.) and a current probe (A621, Tektronix, Inc.), respectively, which were both connected to a digital oscilloscope (TDS1001B, Tektronix, Inc.). The gas flow rate was adjusted using a gas flow meter (RK-1250, Kofloc Kyoto, Kojima Instruments Inc.).

Absorption spectra of the methylene blue aqueous solutions were obtained using a spectrophotometer (V-630, JASCO Corporation). For HPLC, a pump (PU-2080 Plus, JASCO Corporation), a UV detector (UV-2075 Plus, JASCO Corporation), and a column (Hydrosphere C18, YMC Co., Ltd.) were used. A 0.05% aqueous solution of phosphoric acid was used as the mobile phase at a flow rate of 1.0 mL/min. The UV wavelength of the detection unit was 210 nm. The amount of specimen used for the measurement was 20 µL. For emission spectroscopy of the plasma jet, a multichannel spectroscope (QE65000, Ocean Optics) was used. For emission spectroscopy in Ar gas, a quartz glass window was installed on the acrylic resin container to enable the observation of emissions inside the container. The quartz glass window has a transmittance of  $\geq 85\%$  at wavelengths of 190-1100 nm.

Figure 2 shows the circuit used to measure current. To measure the current during plasma jet irradiation, copper foil was used as an electrode to which a 100  $\Omega$  resistor was connected. One end of the resistor was grounded. The

voltage between the two ends of the resistor was measured to obtain the current. In the case of plasma jet irradiation in Ar gas, this circuit was placed in the acrylic resin container.

## 3. Methylene Blue

Figures 3(a) and 3(b) show the molecular structures of methylene blue ( $C_{16}H_{18}ClN_3S$ ) and phenothiazine ( $C_{13}H_9NS$ ), respectively. As the name defined by the International Union of Pure and Applied Chemistry (IUPAC), i.e., 3,7-bis (dimethyl amino) phenothiazin-5-ium chloride, indicates, methylene blue is an organic compound having the basic structure of phenothiazine with side chains of dimethyl amino groups [-N(CH<sub>3</sub>)<sub>2</sub>]. The basic structure is a conjugated system, in which single bonds and double bonds are alternately connected. The  $\pi$  electrons in the conjugated system are delocalized <sup>26)</sup>. The conjugated system of the phenothiazine structure is a chromophore and is the main cause of the color. The side-chain dimethyl amino groups are auxochromes. It is known that the bonding of an auxochrome (side chain) to a chromophore (conjugated system) increases the intensity of the absorption maximum and shifts its wavelength to the longer wavelength side <sup>27)</sup>. This is considered to result from the delocalization of  $\pi$ electrons in the conjugated system, which leads to a decrease in the energy of the  $\pi$  electrons. Negatively charged chlorine ions (Cl<sup>-</sup>) are bonded to positively charged sulfur ions  $(S^+)$  via Coulomb's force. In the methylene blue aqueous solution, methylene blue molecules are positively ionized and  $Cl^{-}$  is the counterion (separated from  $S^{+}$  ions).

### 4. Results and Discussion

Figure 4 shows the plasma jet irradiating the surface of a methylene blue aqueous solution in (a) air and (b) Ar gas. In both cases, in the plasma, there are many streamers that reach the aqueous solution. Therefore, high-energy electrons in the streamers are considered to collide with H<sub>2</sub>O molecules on the surface of the aqueous solution, inducing the dissociation of H<sub>2</sub>O molecules and the generation of OH radicals and hydrogen (H) atoms. The OH radicals are considered to contribute to the decomposition of methylene blue molecules. The dissociation energy of H<sub>2</sub>O molecules into H atoms and OH radicals is reported to be 5.1 eV <sup>28, 29)</sup>. Therefore, the energy of high-energy electrons in a plasma jet is considered to be at least 5.1 eV.

To confirm whether the acrylic resin container is filled with Ar gas instead of air during plasma jet irradiation in Ar gas, emission spectroscopy of the Ar plasma jet was performed. Figure 5 shows the emission spectra of the Ar





(a)



Fig. 4 Plasma jet irradiation in (a) air and (b) Ar gas on the surface of the methylene blue aqueous solution.

plasma jet for irradiation in (a) air and (b) Ar gas. In the spectrum shown in Fig. 5(a), peaks are observed at wavelengths of 300–400 and 700–950 nm. The peak at 309 nm is attributed to the presence of OH radicals  $^{30-32)}$ , peaks at 310–400 nm are attributed to excited N<sub>2</sub> molecules of the 2nd positive system  $^{33-35)}$ , and peaks at 700–950 nm are attributed to excited Ar atoms  $^{36-38)}$ . In contrast, in the spectrum shown in Fig. 5(b), the peaks at 310–400 nm attributed to N<sub>2</sub>



Fig. 5 Emission spectra of plasma jet for irradiation in (a) air and (b) Ar gas.



Fig. 6 Changes in color of methylene blue aqueous solutions irradiated with plasma jet in air (top) and in Ar gas (bottom): (a) before irradiation, (b) after 1 min of irradiation, (c) after 3 min of irradiation, (d) after 5 min of irradiation,(e) after 10 min of irradiation, and (f) after 20 min of irradiation.

molecules are not observed. This result indicates that there are no  $N_2$  molecules in the acrylic resin container and thus no interaction between the plasma and  $N_2$  molecules.

Therefore, it was confirmed that the air in the acrylic resin container was almost completely replaced with Ar gas. The maximum emission intensity of the excited Ar atoms for irradiation in Ar gas was ~60000, which is approximately fourfold higher than that for irradiation in air (~15000). This finding indicates that the number of excited Ar atoms is higher when irradiation is carried out in Ar gas. In other words, the number of ionized Ar atoms is high and the quantity of electrons is also high when irradiation is carried out in Ar gas.

To compare the quantity of electrons that reached the surface of the methylene blue aqueous solutions, the current during plasma jet irradiation in air and in Ar gas was measured using the circuit shown in Fig. 2. The results of the measurements indicate that the current for irradiation in Ar gas is 60.8 mA, which is approximately 1.6-fold higher than that in air (37.6 mA). This finding indicates that the quantity of electrons generated for irradiation in Ar gas is approximately 1.6-fold higher than that in air and that more OH radicals are generated in the methylene blue aqueous solution irradiated with a plasma jet in Ar gas. Therefore, it is expected that decolorization of the methylene blue aqueous solution irradiated with a plasma jet in Ar gas will proceed faster than that of the solution irradiated with a plasma jet in air.

Figure 6 shows the changes in color of methylene blue aqueous solutions irradiated with a plasma jet in air and in Ar gas. Before irradiation, the color of the solution was blue in both cases. With increasing plasma irradiation time, the color gradually changed to lighter blue. The solutions became colorless after 20 and 10 min of irradiation in air and in Ar gas, respectively. The results of emission spectroscopy and current measurements demonstrate that the decolorization of the methylene blue aqueous solution irradiated with a plasma jet in Ar gas would proceed faster than that of the solution irradiated with a plasma jet in air, as we predicted.

The following redox reaction of methylene blue is known: when protons associate with methylene blue molecules in a methylene blue aqueous solution (i.e., methylene blue is methylene reduced), blue becomes the colorless leucomethylene blue. In contrast, when protons are dissociated from leucomethylene blue (i.e., methylene blue is oxidized), leucomethylene blue becomes methylene blue <sup>39)</sup>. In this study, when the methylene blue aqueous solution that was decolored by plasma jet irradiation was stirred, bringing it into contact with air for oxidation, the color of the solution did not return to blue. This finding indicates that the decolorization of the methylene blue aqueous solution upon plasma jet irradiation is not related to the leucomethylene

blue obtained by the reduction of methylene blue. Therefore, the decolorization of the methylene blue aqueous solution irradiated with a plasma jet is related to the decomposition of methylene blue molecules.

Figure 7 shows absorption spectra of methylene blue aqueous solutions irradiated with a plasma jet in (a) air and (b) Ar gas. In the absorption spectra obtained before irradiation, peaks were observed at wavelengths of 246, 291, 614, and 664 nm. In both Figs. 7(a) and 7(b), the absorbance at these peaks decreased with increasing plasma irradiation time. The decrease in absorbance indicates a decrease in the methylene blue concentration because the absorbance is proportional to the methylene blue concentration. This also indicates that the phenothiazine structure is decomposed because the phenothiazine structure in methylene blue is a chromophore.

In our previous studies, we reported that the decomposition of methylene blue molecules is caused by the dissociation of the bonds of the N and sulfur (S) atoms in the phenothiazine structure in the methylene blue aqueous solution irradiated with an Ar plasma jet, on the basis of the results of NMR and ESI-MS<sup>22)</sup>.

In the absorption spectra obtained after irradiation in air, a new peak at a wavelength of approximately 200 nm was observed [Fig. 7(a)]. We have predicted that a new peak should be generated at a wavelength of approximately 200 nm in the absorption spectra of the methylene blue aqueous solution irradiated with a plasma jet in air, similar to the case of plasma jet irradiation onto distilled water, as explained in the Introduction. This prediction was demonstrated to be correct from our findings. This peak is attributed to the presence of NO3<sup>-</sup>, NO2<sup>-</sup>, and H2O2. In contrast, in the absorption spectra obtained after irradiation in Ar gas, a small peak was observed at a wavelength of approximately 200 nm, as shown in Fig. 7(b). The plasma did not come into contact with air; therefore, the peak is not attributed to the presence of  $NO_3^-$  or  $NO_2^-$ ; rather, it is attributed to the presence of H<sub>2</sub>O<sub>2</sub>. Absorption spectra similar to those in Fig. 7(b) are also obtained by underwater discharge inside bubbles in a methylene blue aqueous solution <sup>15, 17)</sup>.

Figure 8 shows the dependence of absorbance at a wavelength of 664 nm on plasma irradiation time. The absorbance at 664 nm is proportional to the methylene blue concentration. The methylene blue concentration in the solution decreases with increasing plasma irradiation time. The absorbance becomes almost zero after 20 min of irradiation in air and 10 min of irradiation in Ar gas. Therefore, the time required for decolorization of the methylene blue aqueous solution irradiated with the plasma jet in Ar gas was half that required for irradiation in air. A



Fig. 7 Absorption spectra of methylene blue aqueous solutions irradiated with plasma jet in (a) air and (b) Ar gas.



Fig. 8 Dependence of absorbance at wavelength of 664 nm on plasma irradiation time.



Fig. 9 Dependence of absorbance at wavelength of 210 nm on plasma irradiation time.



Fig. 10 HPLC chromatograms of methylene blue aqueous solution: (a) before irradiation, and irradiated with plasma jet for 1 min in (b) air and (c) Ar gas.

decrease in the methylene blue concentration with increasing plasma irradiation time has also been reported for atmospheric-pressure plasma irradiation on the surface of a methylene blue aqueous solution <sup>7)</sup> and for plasma discharge inside bubbles in a methylene blue aqueous solution <sup>16)</sup>.

Figure 9 shows the dependence of absorbance on plasma irradiation time at a wavelength of 210 nm. The wavelength of 210 nm matches the wavelength used in HPLC measurement discussed later. The absorbance at this wavelength is considered to be linear with the total amount of  $NO_3^- + NO_2^- + H_2O_2$ . The absorbance at this wavelength increases linearly for 10 min of plasma irradiation, after

which it saturates. In contrast, the absorbance negligibly increases with plasma irradiation in Ar gas. This finding indicates that little  $NO_3^-$ ,  $NO_2^-$ , or  $H_2O_2$  is generated upon plasma irradiation in Ar gas.

Figure 10 shows HPLC chromatograms of methylene blue aqueous solutions irradiated with a plasma jet for 1 min. As shown in Fig. 10(a), no peaks were observed in the chromatogram before irradiation. In the chromatogram of themethylene blue aqueous solution irradiated with a plasma jet in air [Fig. 10(b)], peaks were observed at retention times of 3.0, 4.1, and 5.7 min. From the results of our previous study, these peaks were identified as  $H_2O_2$ ,  $NO_2^-$ , and  $NO_3^-$ , respectively  $^{24)}$ . From the findings of this study,  $H_2O_2$ ,  $NO_2^-$ , and  $NO_3^-$  were confirmed to be generated in the methylene blue aqueous solution irradiated with a plasma jet in air. From the results of comparisons of peak intensity, the proportion of the absorbance of NO<sub>3</sub><sup>-</sup> among the total absorbance at 210 nm was the highest, followed by those of  $H_2O_2$  and  $NO_2^-$  in that order. Most of the absorbance was attributable to NO<sub>3</sub><sup>-</sup>. In the chromatogram of the methylene blue aqueous solution irradiated with a plasma jet in Ar gas [Fig. 10(c)], peaks were observed at retention times of 3.0 and 5.7 min; however, no peak was observed at 4.1 min. This finding indicates that in the methylene blue aqueous solution irradiated with plasma jet in Ar gas, H<sub>2</sub>O<sub>2</sub> and a small amount of NO3<sup>-</sup> were generated; however, no NO2<sup>-</sup> was generated.

Figure 11 shows the dependence of the  $H_2O_2$ concentration obtained using HPLC in the methylene blue aqueous solution irradiated with a plasma jet on plasma irradiation time. The H<sub>2</sub>O<sub>2</sub> concentration was determined from the intensity of the H<sub>2</sub>O<sub>2</sub> peak in chromatograms using a calibration curve. The H<sub>2</sub>O<sub>2</sub> concentration increased linearly with increasing plasma irradiation time for irradiation in air and in Ar gas. The H<sub>2</sub>O<sub>2</sub> concentration in methylene blue aqueous solution after 20 min of irradiation in Ar gas was 70 mg/L, which is ~1.4-fold higher than that in air (51 mg/L). This finding suggests that ~1.4-fold more OH radicals were generated in the solution irradiated with a plasma jet in Ar gas than that in air. This value is close to the ~1.6-fold higher current flowing during plasma jet irradiation in Ar gas compared with that in air. Therefore, we suggest that high-energy electrons in plasma were mostly used for the generation of OH radicals in solutions irradiated with a plasma jet in Ar gas. Similarly, an increase in the H<sub>2</sub>O<sub>2</sub> concentration with increasing plasma irradiation time time was reported and discharge for surface atmospheric-pressure plasma irradiation <sup>40, 41)</sup> and underwater pulse discharge <sup>12, 31, 42)</sup>.

Figure 12 shows the dependences of the NO<sub>3</sub><sup>-</sup>



Fig. 11 Dependence of  $H_2O_2$  concentration obtained by HPLC in methylene blue aqueous solutions irradiated with plasma jet on plasma irradiation time.



Fig. 12 Dependence of  $NO_3^-$  concentration obtained by HPLC in methylene blue aqueous solutions irradiated with plasma jet on plasma irradiation time.

concentration obtained by HPLC in methylene blue aqueous solutions irradiated with a plasma jet on plasma irradiation time. The  $NO_3^-$  concentration was obtained from the intensity of the  $NO_3^-$  peak in chromatograms using a calibration curve. The  $NO_3^-$  concentration increased linearly with increasing plasma irradiation time for irradiation in air; however, it rarely increased for irradiation in Ar gas. This finding indicates that the plasma did not come into contact with  $N_2$  or  $O_2$  molecules in Ar gas owing to the absence of air.

On the basis of the results from ion chromatography, Hunag *et al.* reported that  $NO_3^-$  is generated in pure water irradiated with a dielectric barrier discharge in air and that the  $NO_3^-$  concentration increases with increasing discharge time <sup>9)</sup>. Also, on the basis of the results of ion chromatography, Ikawa *et al.* reported that  $NO_3^-$  and  $NO_2^$ are generated in distilled water irradiated with an atmospheric-pressure helium (He) plasma jet and that the  $NO_3^- + NO_2^-$  concentration linearly increases with increasing plasma irradiation time <sup>43)</sup>. Our results are in good agreement with their results.

The NO<sub>3</sub><sup>-</sup> concentrations in the methylene blue aqueous solutions after 20 min of plasma irradiation in Ar gas and in air were 1.2 and 63.3 mg/L, respectively. Very little NO<sub>3</sub><sup>-</sup> was generated in the solution irradiated with a plasma jet in Ar gas, approximately  $\sim 2\%$  of the amount generated in air.

## 5. Conclusions

Decolorization of methylene blue aqueous solutions irradiated with an atmospheric-pressure Ar plasma jet in air was compared with that in Ar gas. The time required for decolorization of the methylene blue aqueous solution irradiated with a plasma jet in Ar gas was 10 min, which was half that required in air (20 min) because the current for irradiation in Ar gas was 60.8 mA, which was approximately 1.6-fold higher than that in air (37.6 mA). In addition, in the case of plasma jet irradiation in Ar gas, neither NO<sub>3</sub><sup>-</sup> nor NO<sub>2</sub><sup>-</sup> was generated and only H<sub>2</sub>O<sub>2</sub> was generated in the methylene blue aqueous solution, unlike the case of plasma jet irradiation in air. The  $H_2O_2$  concentration in the methylene blue aqueous solution after 20 min of irradiation in Ar gas was 70 mg/L, which was ~1.4-fold higher than that in air (51 mg/L).

The amount of OH radicals generated is large with plasma irradiation in Ar gas because the number of high-energy electrons that collide with  $H_2O$  molecules on the surface of the aqueous solution is high. The reason for the halving of the time required for the decolorization of methylene blue aqueous solution is considered to be the presence of a large amount of OH radicals that contribute to the degradation of methylene blue molecules.

In this study, the irradiation distance was 15 mm. The irradiation distance is considered to be a parameter that limits decolorization. We will carry out experiments with various irradiation distances in Ar gas in the future.

On the basis the results obtained in this study, we hypothesize that harmful organic compounds in wastewater can be decomposed in a shorter time without generating  $NO_x^-$  by atmospheric-pressure Ar plasma jet irradiation in Ar gas than that in air.

#### References

 B. Sun, M. Sato and J. S. Clements: Use of a Pulsed High-Voltage Discharge for Removal of Organic Compounds in Aqueous Solution, J. Phys. D: Appl. Phys., 32, 1908 (1999).

- W. F. L. M. Hoeben, E. M. van Veldhuizen, W. R. Rutgers, and G. M. W. Kroesen: Gas Phase Corona Discharges for Oxidation of Phenol in an Aqueous Solution, J. Phys. D: Appl. Phys. 32, L133 (1999).
- D. Hayashi, W. F. L. M. Hoeben, G. Dooms, E. M. van Veldhuizen, W. R. Rutgers, and G. M. W. Kroesen: Influence of Gaseous Atmosphere on Corona-induced Degradation of Aqueous Phenol, J. Phys. D: Appl. Phys. 33, 2769 (2000).
- L. R. Grabowski, E. M. van Veldhuizen, A. J. M. Pemen, and W. R. Rutgers: Breakdown of Methylene Blue and Methyl Orange by Pulsed Corona Discharge, Plasma Sources Sci. Technolo. 16, 226 (2007).
- S. Ikoma, K. Satoh, and H. Itoh: Decomposition of Methylen Blue in an Aqueous Solution using a Pulsed-discharge Plasma at Atmospheric-Pressure, IEEJ Trans. FM, 129, 237 (2009) (in Japanese).
- T. Miichi: Decolorization of Indigo Carmine Solution using Discharge on Surface of a Gas-layer in Water, IEEJ Trans. FM, **126**, 851 (2006) (in Japanese).
- K. Kitano, H. Aoki, and S. Hamaguchi: Radio-Frequency-Driven Atmospheric-Pressure Plasmas in Contact with Liquid Water, Jpn. J. Appl. Phys., 45, 8294 (2006).
- G. Chen, M. Zhou, S. Chen, and W. Chen: The Different Effects of Oxygen and Air DBD Plasma by Products on the Degradation of Methyl Violet 5BN, J. Haz. Mater. 172, 786 (2009).
- F. Huang, L. Chen, H. Wang, and Z. Yan: Analysis of the Degradation Mechanism of Methylene Blue by Atmospheric Pressure Dielectric Barrier Discharge Plasma, Chem. Eng. J., 162, 250 (2010).
- Wahyudiono, S. Machmudah, and M. Goto: Pulsed Discharge Plasma over a Water Surface Induces Decoloration of Dyes, J. Phys.: Conference Series 441, 012008 (2013).
- J. S. Clements, M. Sato, and R. H. Davis: Preliminary Investigation of Prebreakdown Phenomena and Chemical Reactions using a Pulsed High-voltage Discharge in Water, IEEE Trans. Ind. Appl. 23, 224 (1987).
- 12) A. Joshi, B. R. Locke, P. Arce, and W. C. Finney: Formation of Hydroxyl Radicals, Hydrogen Peroxide and Aqueous Electrons by Pulsed Streamer Corona Discharge in Aqueous Solution, J. Haz. Mater. 41, 3 (1995).
- A. T. Sugiarto, S. Ito, T. Oshima, M. Sato, and J. D. Skalny: Oxidative Decoloration of Dyes by Pulsed Discharge Plasma in Water, J. Electrostat. 58, 135 (2003).

- 14) K. Sharama, B. R. Locke, P. Arce and W. C. Finney: A Preliminary Study of Pulsed Streamer Corona Discharge for the Degradation of Phenol in Aqueous Solutions, Hazard Waste Hazard Mater. 10, 209 (1993).
- T. Maehara, H. Toyota, M. Kuramoto, A. Iwamae, A. Tadokoro, S. Mukasa, H. Yamashita, A. Kawashima and S. Nonura: Radio Frequency Plasma in Water, Jpn. J. Appl. Phys. 45, 8864 (2006).
- 16) T. Ishijima, H. Hotta, H. Sugai, and M. Sato: Multibubble Plasma Production and Solvent Decomposition in Water by Slot-Excited Microwave Discharge, Appl. Phys. Lett., 91, 121501 (2007).
- 17) I. Miyamoto, T. Maehara, H. Miyaoka, S. Onishi, S. Mukasa, H. Toyota, M. Kuramoto, S. Nomura, and A. Kawashima: Effect of the Temperature of Water on the Degradation of Methylene Blue by the Generation of Radio Frequency Plasma in Water, J. Plasma Fusion Res. SERIES, 8, 627 (2009).
- W. M. Latimer: The Oxidation States of the Elements and Their Potentials in Aqueous Solutions, 2nd. Ed., Prentice-Hall, p. 48 (1952).
- B. Sun, M. Sato, and J. S. Clements: Optical Study of Active Species Produced by a Pulsed Streamer Corona Dischsrge in Water, J. Electrostatics 39, 189 (1997).
- 20) M. A. Malik, A. Ghaffar, and S. A. Malik: Water Purification by Electrical Discharges, Plasma Source Sci. Technol., 10, 82 (2001).
- 21) H. Kuwahata, K. Kimura, and R. Ohyama: Decolorization of Methylene Blue Aqueous Solution by Atmospheric-Pressure Plasma Jet, e-J. Surf. Sci. Nanotech. 8, 381 (2010).
- 22) M. Uchida, S. Kezuka, Y. Oda, M. Isomura, and H. Kuwahata: Decolorization of Methylene Blue Aqueous Solution by Atmospheric-pressure Plasma Jet Irradiation: NMR and MS Analyses, Proc. Sch. Eng. Tokai Univ., Ser. J, Vol. 56, No. 2, 9 (2016) (in Japanese).
- 23) H. Kuwahata, K. Kimura, and I. Mikami: Identification of Peak near 200 nm in Absorbancw Spectrum of Distilled Water upon Atmospheric-Pressure Plasma Jet Irradiation, e-J. Surf. Sci. Nanotech., 9, 442 (2011).
- 24) H. Kuwahata and I. Mikami: High Performance Liquid Chromatography Analysis of Distilled Water Irradiated with Atmospheric-pressure Plasma Jet, Proc. Sch. Eng. Tokai Univ., Ser. E, 40, 1 (2015).
- 25) R. Ohyama and A. Nagai: Japan Patent 244938 (2006).
- J. March: Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, 2<sup>nd</sup> Ed., McGraw-Hill p. 32 (1977).
- 27) J. R. Dyer: Applications of Absorption Spectroscopy of

Organic Compounds, Prentice Hall Inc. p. 19 (1965).

- 28) Y. R. Luo: *Comprehensive Handbook of Chemical Bond Energies*, CRC Press, Boca Raton, FL, (2007).
- 29) V. I. Vedeneyev, L. V. Gurvich, V. N. Kondrat'yev, V. A. Medvedev, and Ye. L. Frankevich: *Bond Energies, Ionization Potentials and Electron Affinities*, St. Martin's Press, New York, (1962).
- 30) H. Okabe: Photodissociation of Nitric Acid and Water in The Vacuum Ultraviolet; Vibrational and Rotational Distributions of  $OH^2\Sigma^+$ , J. Chem. Phys. **72**, 6642 (1980).
- M. Sato, T. Ohgiyama, and J. S. Clements: Formation of Chemical Species and their Effects on Microorganisms Using a Pulsed High-voltage Discharge in Water, IEEE Trans. Ind. Appl., 32, 106 (1996).
- 32) C. Sugama, F. Tochikubo, and S. Uchida: Glow Discharge Fomation over Water Surface at Saturated Water Vapor Pressure and Its Application to Wastewater Treatment, Jpn. J. Appl. Phys. 45, 8858 (2006).
- 33) A. Lofthus and P. H. Krupenie: The Spectrum of Molecular Nitrogen, J. Phys. Chem. Ref. Date, Vol. 6, No. 1, 113 (1977).
- 34) T. Yuji, T. Urayama, S. Fujii, Y. Iijima, Y. Suzaki, and H. Akatsuka: Basic Characteristics for PEN Film Surface Modification using Atmospheric-Pressure Non-equilibrium Microwave Plasma Jet, IEEJ Trans. FM, 128, 449 (2008) (in Japanese).
- 35) S. J. Kim, T. H. Chung, S. H. Bae, and S. H. Leem: Characterization of Atmospheric Pressure Microplasma Jet Source and its Application to Bacterial Inactivation, Plasma Process. Polym. 6, 676 (2009).

- G. R. Harrison: *MIT Wavelength Tables*, MIT Press, Cambridge (1969).
- 37) H. Kuwahata and R. Ohyama: Fundamental Characteristics of Atmospheric Pressure Argon Plasma Jet and Surface Modification of Glasses, Proc. Sch. Eng. Tokai Univ., Ser. J, Vol. 48, No. 2, 17 (2008) (in Japanese).
- 38) D. Shibahara, H. Yasuda, N. Fujii, H. Kurita, K. Takashima, and A. Mizuno: Sterization of Escherichia coli by Small Size Argon Plasma Jet, J. Inst. Electrostat. Jpn. 34, 2 (2010) (in Japanese).
- S. Mowry and P. J. Ogren: Kinetics of Methylene Blue Reduction by Ascorbic Acid, J. Chem. Educ. 76, 970 (1999).
- K. Oehmigen, M. Hahnel, R. Brandenburg, Ch. Wilke, K. -D. Weltmann, and Th. von Woedtke: The Role of Acidification for Antimicrobial Activity of Atmospheric Pressure Plasma in Liquids, Plasma Process. Polym. 7, 250 (2010).
- 41) C. A. J. van Gils, S. Hofmann, B. K. H. L. Boekema, R. Brandenburg, and P. J. Bruggeman: Mechanisms of Bacterial Inactivation in the Liquid Phase Induced by a Remote RF Cold Atmospheric Pressure Plasma Jet, J. Phys. D: Appl. Phys. 46, 175203 (2013).
- 42) Ruma, N. Aoki, T. Sakugawa, H. Akiyama, and M. Akiyama: Hydrogen Peroxide Generation by Pulsed Discharge in Bubbling Water, IEEJ Trans. FM, 133, 636 (2013).
- 43) S. Ikawa, K. Kitano, and S. Hamaguchi: Effects of pH on Bacterial Inactivation in Aqueous Solutions due to Low-Temperature Atmospheric Pressure Plasma Application, Plasma Process. Polym. 7, 33 (2010).